

Recursion Rules for The Hypergeometric Zeta Function

Lin Jiu

Tulane University

Joint Work with Alyssa Byrnes, Victor H. Moll and Christophe Vignat

Supported by NSF-DMS 0070567

Outlines

1 Motivation & Background

- Zeta Functions
- Bernoulli Numbers

2 Hypergeometric Zeta Function

- Kummer Hypergeometric Function
- Hypergeometric Zeta Function
- Main Results

3 A Probabilistic Approach

- Moments & Cumulants
- Conjugate Random Variables

Question

Definition

$$\Phi_{a,b}(z) := M(a, a+b, z) = {}_1F_1\left(\begin{matrix} a \\ a+b \end{matrix} \middle| z\right) = \sum_{n=0}^{\infty} \frac{(a)_n}{(a+b)_n} \cdot \frac{z^n}{n!},$$

where

$$(a)_n := a(a+1)\cdots(a+n-1)$$

is the Pochhammer symbol.

Objects

The sequence $\{A_{a,b}\}$ defined by the generating function

$$\frac{1}{\Phi_{a,b}(z)} = \sum_{n=0}^{\infty} A_{a,b} \frac{z^n}{n!}.$$



Question (Continued)

Objects

The roots of $\Phi_{a,b}(z)$, denoted by

$$\{z_{a,b;n}^s : n \in \mathbb{N}\}$$

and the corresponding hypergeometric-zeta function

$$\zeta_{a,b}^H(s) = \sum_{n=0}^{\infty} \frac{1}{z_{a,b;n}^s}, .$$

Zeta Functions

Example

Riemann-Zeta Function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \operatorname{Re}(s) > 1.$$

$\mathbb{A} := \{1, 2, \dots\} = \{a_n\}_{n \in \mathbb{N}}$, where $a_n = n$

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{a_n^s}.$$

$$f(z) := \frac{\sin(\pi z)}{\pi z} \implies \mathbb{A} = \{z \in \mathbb{C} : f(z) = 0, z > 0\}.$$

Zeta Functions (Continued)

Example

Bessel-Zeta Function

$$\zeta_{Bes,a}(s) := \sum_{n=1}^{\infty} \frac{1}{j_{a,n}^s},$$

where $\mathbb{A} := \{j_{a,n}\}$ are the zeros of $J_a(z)/z^a$ for the Bessel function of the first kind

$$J_a(z) := \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m+a+1)} \left(\frac{z}{2}\right)^{2m+a}.$$

Zeta Functions (Continued)

In general, given (/selected) function f , we could define

$$\{a_n\} := \mathbb{A}(f) := \{z \in \mathbb{C} : z \neq 0, f(z) = 0\}$$

and

$$\zeta_f(s) = \zeta_{\mathbb{A}}(s) = \sum_{n=1}^{\infty} \frac{1}{a_n^s}.$$

Bernoulli Numbers

Definition

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!} \text{ for } |x| < 2\pi.$$

Theorem

$$\zeta(2n) = (-1)^{n+1} \frac{(2\pi)^{2n}}{2(2n)!} B_{2n}.$$

Bernoulli Numbers (Continued)

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}$$

L. Carlitz 1961

$$\frac{x^2}{e^x - 1 - x} = \sum_{n=0}^{\infty} \beta_n \frac{x^n}{n!}.$$

Kummer Hypergeometric Function

Definition

$$\Phi_{a,b}(z) := M(a, a+b, z) = {}_1F_1 \left(\begin{matrix} a \\ a+b \end{matrix} \middle| z \right) = \sum_{n=0}^{\infty} \frac{(a)_n}{(a+b)_n} \cdot \frac{z^n}{n!},$$

where

$$(a)_n := a(a+1)\cdots(a+n-1)$$

is the Pochhammer symbol.

Fact

$$\frac{\frac{x^k}{k!}}{e^x - \sum_{s=0}^{k-1} \frac{x^s}{s!}} = \frac{1}{\Phi_{1,k}(x)}.$$



Hypergeometric Zeta Function

Definition

The Hypergeometric Zeta Function is defined by

$$\zeta_{a,b}^H(s) = \sum_{n=0}^{\infty} \frac{1}{z_{a,b;n}^s},$$

where $\mathbb{A} := \{z_{a,b;n} : n \in \mathbb{N}\}$ is the set of zeros of $\Phi_{a,b}(z)$.

Connection

Fact

{A. Byrnes, -, V. Moll, C. Vignat} *By applying Hadamard Factorization Theorem,*

$$\Phi_{a,b}(z) = e^{\frac{a}{a+b}z} \prod_{k=1}^{\infty} \left(1 - \frac{z}{z_{a,b;k}}\right) e^{z/z_{a,b;k}}.$$

{A. Byrnes, -, V. Moll, C. Vignat} *By Contiguous Relation*

$$\frac{\Phi_{a,b+1}(z)}{\Phi_{a,b}(z)} = 1 + \frac{a+b}{b} \sum_{k=1}^{\infty} \zeta_{a,b}^H(k+1) z^k.$$

Recurrences

Theorem

{A. Byrnes, -, V. Moll, C. Vignat} [Linear Recurrence]

$$\sum_{k=1}^p B(a+p+k, b) \frac{p!}{(p-k)!} \zeta_{a,b}^H(k+1) = -\frac{bp}{(a+b)(a+b+p)} B(a+p, b),$$

where

$$B(u, v) := \int_0^1 x^{u-1} (1-x)^{v-1} dx$$

is the Beta function.

Recurrences (Continued)

Theorem

{A. Byrnes, -, V. Moll, C. Vignat} [Quadratic Recurrence]

$$\sum_{k=1}^p \zeta_{a,b}^H(k+1) \zeta_{a,b}^H(p-k+1) = (a+b+p+1) \zeta_{a,b}^H(p+2) + \left(\frac{a-b}{a+b} \right) \zeta_{a,b}^H(p+1),$$

which is the analogue of

$$(n+1) \zeta(2n) = \sum_{k=1}^{n-1} \zeta(2k) \zeta(2n-2k), \text{ for } n \geq 2.$$

Hypergeometric Bernoulli Numbers

Definitions

$$\frac{1}{\Phi_{1,b}(z)} = \sum_{n=0}^{\infty} B_n^{(b)} \frac{z^n}{n!}.$$

Theorem

{A. Byrnes, -, V. Moll, C. Vignat}

$$B_n^{(b)} = \begin{cases} 1 & n = 0 \\ -\frac{1}{b+1} & n = 1 \\ -\frac{n!}{b} \zeta_{1,b}^H(n) & n \geq 2 \end{cases}.$$

Remark

It suggests

$$\zeta_{1,b}^H(1) = \frac{b}{1+b}.$$



A Probabilistic Approach

Moments & Cumulants

For a given random variable X with continuous density function $r(x)$,

(I) the expectation operator is defined by

$$\mathbb{E}u(X) := \int_{\mathbb{R}} u(x) r(x) dx;$$

(II) the moments are $\mathbb{E}[X^n]$ and the moment generating function is given by

$$\varphi_X(t) := \mathbb{E}[e^{tX}] = \int_{\mathbb{R}} e^{tx} r(x) dx = \sum_{n=0}^{\infty} \mathbb{E}[X^n] \frac{t^n}{n!};$$

(III) the cumulants $\kappa_X(n)$ are given by

$$\log \varphi_X(t) = \sum_{m=1}^{\infty} \kappa_X(m) \frac{z^m}{m!}.$$

Moments & Cumulants (Continued)

Example

[Exponential Distribution] Random variable Γ has density function

$$f_{\Gamma}(x) = \begin{cases} e^{-x} & x \geq 0 \\ 0 & x < 0 \end{cases} \implies \mathbb{E}[\Gamma^a] = \Gamma(a+1),$$

then

$$\mathbb{E}[e^{t\Gamma}] = \frac{1}{1-t}, \text{ for } |t| < 1.$$

Moments & Cumulants (Continued)

Example

[Beta Distribution] Random variable $\mathcal{B}_{a,b}$ has density function

$$f_{\mathcal{B}_{a,b}}(x) = \begin{cases} \frac{x^{a-1}(1-x)^{b-1}}{B(a,b)} & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

then

$$\mathbb{E}[e^{t\mathcal{B}_{a,b}}] = \frac{1}{B(a,b)} \int_0^1 e^{tx} x^{a-1} (1-x)^{b-1} dx = \Phi_{a,b}(t).$$

Moments & Cumulants (Continued)

Theorem

For a general random variable X ,

$$\kappa_X(n) = \mathbb{E}[X^n] - \sum_{j=1}^{n-1} \binom{n-1}{j-1} \kappa_X(j) \mathbb{E}[X^{j-1}].$$

{A. Byrnes, -, V. Moll, C. Vignat}

$$(n-1)! \sum_{k=2}^n \frac{B(a+n-k, b)}{(n-k)!} \zeta_{a,b}^H(k) = \frac{a}{a+b} B(a+n-k, b) - B(a+n, b),$$

which is just the linear recurrence mentioned before.

Moments & Cumulants (Continued)

Theorem

{A. Byrnes, -, V. Moll, C. Vignat}

$$(n-1)! \sum_{k=2}^n \frac{B(a+n-k, b)}{(n-k)!} \zeta_{a,b}^H(k) = \frac{a}{a+b} B(a+n-k, b) - B(a+n, b) . (*)$$

Recall

{A. Byrnes, -, V. Moll, C. Vignat} [Linear Recurrence]

$$\sum_{k=1}^p B(a+p+k, b) \frac{p!}{(p-k)!} \zeta_{a,b}^H(k+1) = -\frac{bp}{(a+b)(a+b+p)} B(a+p, b)$$

Conjugate Random Variables

Definition

Two random variables X and Y are called conjugate random variables if

$$\mathbb{E}[(X + Y)^n] = \delta_n = \begin{cases} 1 & n = 0 \\ 0 & \text{otherwise} \end{cases}.$$

Equivalently,

$$1 = \mathbb{E}[e^{t(X+Y)}] = \mathbb{E}[e^{tX}] \mathbb{E}[e^{tY}].$$

Conjugate Random Variables (Continued)

Recall

$$\Phi_{a,b}(z) = e^{\frac{a}{a+b}z} \prod_{k=1}^{\infty} \left(1 - \frac{z}{z_{a,b;k}}\right) e^{z/z_{a,b;k}}, \mathbb{E}[e^{t\Gamma}] = \frac{1}{1-z}$$

Theorem

{A. Byrnes, -, V. Moll, C. Vignat}

$$\mathcal{Z}_{a,b} := -\frac{a}{a+b} + \sum_{k=1}^{\infty} \frac{\Gamma_k - 1}{z_{a,b;k}},$$

where $\{\Gamma_k\}_{k=0}^{\infty}$ is a sequence of i.i.d of exponential distributions, is the conjugate of $\mathcal{B}_{a,b}$, namely

$$\mathbb{E}[e^{t\mathcal{Z}_{a,b}}] = 1/\Phi_{a,b}(t).$$

Conjugate Random Variables (Continued)

Theorem

{A. Byrnes, -, V. Moll, C. Vignat} Let X and Y be conjugate random variables and define polynomials

$$P_n(z) = \mathbb{E}[(z + X)^n] \text{ and } Q_n(z) = \mathbb{E}[(z + Y)^n],$$

then

$$\begin{cases} P_{n+1}(z) - zP_n(z) = \sum_{j=0}^n \binom{n}{j} \kappa_X(j+1) P_{n-j}(z) \\ Q_{n+1}(z) - zQ_n(z) = -\sum_{j=0}^n \binom{n}{j} \kappa_X(j+1) Q_{n-j}(z) \end{cases}$$

Conjugate Random Variables (Continued)

Theorem

{A. Byrnes, -, V. Moll, C. Vignat} $[X = \mathcal{B}_{a,b} \ Y = \mathcal{Z}_{a,b}]$

$$(n-1)! \sum_{j=2}^n \frac{B_{n-j}^{(a,b)}}{(n-j)} \zeta_{a,b}^H(j) = \frac{a}{a+b} B_{n-1}^{(a,b)} + B_n^{(a,b)}, \ (**)$$

where $B_n^{(a,b)}$ are the generalized Bernoulli numbers defined by

$$\sum_{n=0}^{\infty} B_n^{(a,b)} \frac{z^n}{n!} = \frac{1}{\Phi_{a,b}(z)}.$$

Conjugate Random Variables (Continued)

Remark

Identity (**) is different from the linear recurrence obtained through moments and cumulants (*).

$$\begin{cases} (n-1)! \sum_{k=2}^n \frac{B(a+n-k, b)}{(n-k)!} \zeta_{a,b}^H(k) = \frac{a}{a+b} B(a+n-k, b) - B(a+n, b). & (*) \\ (n-1)! \sum_{j=2}^n \frac{B_{n-j}^{(a,b)}}{(n-j)} \zeta_{a,b}^H(j) = \frac{a}{a+b} B_{n-1}^{(a,b)} + B_n^{(a,b)} & (**) \end{cases}$$

Example

Let $a = 5$, $b = 3$ and $n = 3$,

$$\begin{cases} 2\zeta_{5,3}^H(3) + \frac{5}{4}\zeta_{5,3}^H(2) + \frac{1}{32} = 0 \\ 2\zeta_{5,3}^H(3) - \frac{5}{4}\zeta_{5,3}^H(2) - \frac{13}{384} = 0 \end{cases}$$

Conjugate Random Variables (Continued)

Remark

We also have some results on generalized Bernoulli polynomials defined by

$$\sum_{n=0}^{\infty} B_n^{(a,b)}(x) \frac{z^n}{n!} = \frac{e^{xz}}{\Phi_{a,b}(z)}.$$

These polynomials satisfy

$$\sum_{k=0}^n \binom{a+b+n-1}{k} \binom{a+n-k-1}{a-1} B_k^{(a,b)}(x) = (a+b)_n \frac{x^n}{n!}.$$

Future Work

- ① Explanation for two different linear recurrences.
- ② From ${}_1F_1$ to ${}_2F_1$ or even ${}_pF_q$.
- ③ Analogue for “Euler Case”

The End

Thank you for your patience!