

Bernoulli and Euler Symbols: Umbral Calculus, Random Variables, and Multiple Zeta Values

Lin Jiu

DKU-SJTU Workshop

Jan. 5th, 2022

Bernoulli polynomials & Bernoulli numbers

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad \text{and} \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

Bernoulli polynomials & Bernoulli numbers

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad \text{and} \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$



$$\zeta(2n) = \frac{2^{2n}}{2(2n)!} |B_{2n}|$$

Bernoulli polynomials & Bernoulli numbers

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad \text{and} \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$



$$\zeta(2n) = \frac{2^{2n}}{2(2n)!} |B_{2n}| \pi^{2n}$$

Bernoulli polynomials & Bernoulli numbers

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad \text{and} \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$



$$\zeta(2n) = \frac{2^{2n}}{2(2n)!} |B_{2n}| \pi^{2n} \quad \text{and} \quad \zeta(-n) = -\frac{B_{n+1}}{n+1}$$

Bernoulli polynomials & Bernoulli numbers

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad \text{and} \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$



$$\zeta(2n) = \frac{2^{2n}}{2(2n)!} |B_{2n}| \pi^{2n} \quad \text{and} \quad \zeta(-n) = -\frac{B_{n+1}}{n+1} = (-1)^n \frac{B_{n+1}}{n+1}.$$

Bernoulli polynomials & Bernoulli numbers

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad \text{and} \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$



$$\zeta(2n) = \frac{2^{2n}}{2(2n)!} |B_{2n}| \pi^{2n} \quad \text{and} \quad \zeta(-n) = -\frac{B_{n+1}}{n+1} = (-1)^n \frac{B_{n+1}}{n+1}.$$

▶ Euler-Maclaurin Summation Formula

$$\begin{aligned} \sum_{j=a}^n f(j) &= \int_a^n f(x) dx + \frac{f(a) + f(n)}{2} + \sum_{s=1}^m \frac{B_{2s}}{(2s)!} \left(f^{(2s-1)}(n) - f^{(2s-1)}(a) \right) \\ &\quad + \int_a^n \frac{B_{2m} - B_{2m}(\{x\})}{(2m)!} f^{(2m)}(x) dx. \end{aligned}$$

Bernoulli polynomials & Bernoulli numbers

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad \text{and} \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$



$$\zeta(2n) = \frac{2^{2n}}{2(2n)!} |B_{2n}| \pi^{2n} \quad \text{and} \quad \zeta(-n) = -\frac{B_{n+1}}{n+1} = (-1)^n \frac{B_{n+1}}{n+1}.$$

▶ Euler-Maclaurin Summation Formula

$$\begin{aligned} \sum_{j=a}^n f(j) &= \int_a^n f(x) dx + \frac{f(a) + f(n)}{2} + \sum_{s=1}^m \frac{B_{2s}}{(2s)!} \left(f^{(2s-1)}(n) - f^{(2s-1)}(a) \right) \\ &\quad + \int_a^n \frac{B_{2m} - B_{2m}(\{x\})}{(2m)!} f^{(2m)}(x) dx. \end{aligned}$$

▶ Modular forms/Eisenstein series:

$$G_{2k}(\tau) = 2\zeta(2k) \left(1 - \frac{4k}{B_{2k}} \sum_{n=1}^{\infty} \sum_{d|n} d^{2k-1} e^{2\pi i n \tau} \right).$$

Umbral Calculus

$$\mathcal{B}^n = B_n$$

Umbral Calculus

$$\mathcal{B}^n = B_n$$

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k$$

Umbral Calculus

$$\mathcal{B}^n = B_n$$

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k = \sum_{k=0}^n \binom{n}{k} \mathcal{B}^{n-k} x^k$$

Umbral Calculus

$$\mathcal{B}^n = B_n$$

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k = \sum_{k=0}^n \binom{n}{k} \mathcal{B}^{n-k} x^k = (\mathcal{B} + x)^n.$$

Umbral Calculus

$$\mathcal{B}^n = B_n$$

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k = \sum_{k=0}^n \binom{n}{k} \mathcal{B}^{n-k} x^k = (\mathcal{B} + x)^n.$$

$$\triangleright B'_n(x) = nB_{n-1}(x) \Leftrightarrow \frac{d}{dx}(\mathcal{B} + x)^n = n(\mathcal{B} + x)^{n-1}.$$

Umbral Calculus

$$\mathcal{B}^n = B_n$$

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_{n-k} x^k = \sum_{k=0}^n \binom{n}{k} \mathcal{B}^{n-k} x^k = (\mathcal{B} + x)^n.$$

$$\triangleright B'_n(x) = nB_{n-1}(x) \Leftrightarrow \frac{d}{dx}(\mathcal{B} + x)^n = n(\mathcal{B} + x)^{n-1}.$$

Treat $t = \partial_x$, and

$$\frac{t}{e^t - 1} \bullet x^n = B_n(x)$$

Random Variable

$$B_n(x) =$$

Random Variable

$$B_n(x) = \mathbb{E}[(\mathcal{B} + x)^n] =$$

Random Variable

$$B_n(x) = \mathbb{E}[(\mathcal{B} + x)^n] = \mathbb{E}\left[\left(iL - \frac{1}{2} + x\right)^n\right]$$

Random Variable

$$\begin{aligned} B_n(x) &= \mathbb{E}[(\mathcal{B} + x)^n] = \mathbb{E}\left[\left(iL - \frac{1}{2} + x\right)^n\right] \\ &= \frac{\pi}{2} \int_{\mathbb{R}} \left(iL - \frac{1}{2} + x\right)^n \operatorname{sech}^2(\pi t) dt. \end{aligned}$$

Random Variable

$$\begin{aligned} B_n(x) &= \mathbb{E}[(\mathcal{B} + x)^n] = \mathbb{E}\left[\left(iL - \frac{1}{2} + x\right)^n\right] \\ &= \frac{\pi}{2} \int_{\mathbb{R}} \left(iL - \frac{1}{2} + x\right)^n \operatorname{sech}^2(\pi t) dt. \end{aligned}$$

$$\triangleright \left(\frac{t}{e^t - 1}\right)^p e^{tx} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!}$$

Random Variable

$$\begin{aligned} B_n(x) &= \mathbb{E}[(\mathcal{B} + x)^n] = \mathbb{E}\left[\left(iL - \frac{1}{2} + x\right)^n\right] \\ &= \frac{\pi}{2} \int_{\mathbb{R}} \left(iL - \frac{1}{2} + x\right)^n \operatorname{sech}^2(\pi t) dt. \end{aligned}$$

$$\triangleright \left(\frac{t}{e^t - 1}\right)^p e^{tx} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Rightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \cdots + \mathcal{B}_p + x)^n$$

Random Variable

$$\begin{aligned} B_n(x) &= \mathbb{E}[(\mathcal{B} + x)^n] = \mathbb{E}\left[\left(iL - \frac{1}{2} + x\right)^n\right] \\ &= \frac{\pi}{2} \int_{\mathbb{R}} \left(iL - \frac{1}{2} + x\right)^n \operatorname{sech}^2(\pi t) dt. \end{aligned}$$

$$\triangleright \left(\frac{t}{e^t - 1}\right)^p e^{tx} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Rightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \dots + \mathcal{B}_p + x)^n$$

where $(\mathcal{B}_j)_{j=1}^p$ is a sequence of i. i. d. random variables

Random Variable

$$\begin{aligned} B_n(x) &= \mathbb{E}[(\mathcal{B} + x)^n] = \mathbb{E}\left[\left(iL - \frac{1}{2} + x\right)^n\right] \\ &= \frac{\pi}{2} \int_{\mathbb{R}} \left(iL - \frac{1}{2} + x\right)^n \operatorname{sech}^2(\pi t) dt. \end{aligned}$$

$$\triangleright \left(\frac{t}{e^t - 1}\right)^p e^{tx} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Rightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \dots + \mathcal{B}_p + x)^n$$

where $(\mathcal{B}_j)_{j=1}^p$ is a sequence of i. i. d. random variables each

$$\mathcal{B}_j \sim \mathcal{B}.$$

Random Variable

$$\begin{aligned} B_n(x) &= \mathbb{E}[(\mathcal{B} + x)^n] = \mathbb{E}\left[\left(iL - \frac{1}{2} + x\right)^n\right] \\ &= \frac{\pi}{2} \int_{\mathbb{R}} \left(iL - \frac{1}{2} + x\right)^n \operatorname{sech}^2(\pi t) dt. \end{aligned}$$

$$\triangleright \left(\frac{t}{e^t - 1}\right)^p e^{tx} = \sum_{n=0}^{\infty} B_n^{(p)}(x) \frac{t^n}{n!} \Rightarrow B_n^{(p)}(x) = (\mathcal{B}_1 + \dots + \mathcal{B}_p + x)^n$$

where $(\mathcal{B}_j)_{j=1}^p$ is a sequence of i. i. d. random variables each

$$\mathcal{B}_j \sim \mathcal{B}.$$

\triangleright Bernoulli-Barnes Polynomials: let $\mathbf{a} = (a_1, \dots, a_k)$

$$\vec{\mathcal{B}} = (\mathcal{B}_1, \dots, \mathcal{B}_k) \text{ and } |\mathbf{a}| = \prod_{i=1}^k a_i$$

$$e^{tx} \prod_{i=1}^k \frac{t}{e^{a_i t} - 1} = \sum_{n=0}^{\infty} B_n(\mathbf{a}; x) \frac{t^n}{n!} \Leftrightarrow B_n(\mathbf{a}; x) = \frac{1}{|\mathbf{a}|} \left(x + \mathbf{a} \cdot \vec{\mathcal{B}}\right)^n;$$

Theorem (A. Bayad and M. Beck, 2014)

Difference Formula: Suppose $A = \sum_{k=1}^n a_k \neq 0$, then

$$(-1)^m B_m(\mathbf{a}; -x) - B_m(\mathbf{a}; x) = m! \sum_{l=1}^{n-1} \sum_{|L|=l} \frac{B_{m-n+l}(\mathbf{a}_L; x)}{(m-n+l)!},$$

where $L \subset \{1, \dots, n\}$ and $B_m(\mathbf{a}_L; x) = x^m$ if $L = \emptyset$.

Theorem (A. Bayad and M. Beck, 2014)

Difference Formula: Suppose $A = \sum_{k=1}^n a_k \neq 0$, then

$$(-1)^m B_m(\mathbf{a}; -x) - B_m(\mathbf{a}; x) = m! \sum_{l=1}^{n-1} \sum_{|L|=l} \frac{B_{m-n+l}(\mathbf{a}_L; x)}{(m-n+l)!},$$

where $L \subset \{1, \dots, n\}$ and $B_m(\mathbf{a}_L; x) = x^m$ if $L = \emptyset$.

Theorem (L. Jiu, V. Moll and C. Vignat, 2016)

$$f\left(x - \mathbf{a} \cdot \vec{\mathcal{B}}\right) = \sum_{\ell=0}^n \sum_{|L|=\ell} |\mathbf{a}|_{L^*} f^{(n-\ell)}\left(x + \left(\mathbf{a} \cdot \vec{\mathcal{B}}\right)_L\right).$$

$$L^* = \{1, \dots, n\} \setminus L.$$

Corollary

Pick $f(x) = x^m/m!$.

Analytic Continuation: for n_1, \dots, n_r positive integers

Theorem (Sadaoui, 2014)

$$\zeta_r(-n_1, \dots, -n_r) = (-1)^r \sum_{k_2, \dots, k_r} \frac{1}{\bar{n} + r - \bar{k}} \prod_{j=2}^r \frac{\binom{\sum_{i=j}^r n_i - \sum_{i=j+1}^n k_i + r - j + 1}{k_j}}{\sum_{i=j}^r n_i - \sum_{i=j}^n k_i + r - j + 1}$$

$$\times \sum_{l_1, \dots, l_r} \binom{\bar{n} + r - \bar{k}}{l_1} \binom{k_2}{l_2} \cdots \binom{k_r}{l_r} B_{l_1} \cdots B_{l_r},$$

$$\bar{n} = \sum_{j=1}^n n_j, \quad \bar{k} = \sum_{j=2}^r k_j, \quad k_2, \dots, k_r \geq 0, \quad l_j \leq k_j \text{ for } 2 \leq j \leq r \text{ and } l_1 \leq \bar{n} + r + \bar{k}.$$

Theorem (Akiyama and Tanigawa, 2001)

$$\zeta_r(-n_1, \dots, -n_r) = -\frac{\zeta_{r-1}(-n_1, \dots, -n_{r-2}, -n_{r-1} - n_r - 1)}{1 + n_r}$$

$$- \frac{\zeta_{r-1}(-n_1, \dots, -n_{r-2}, -n_{r-1} - n_r)}{2}$$

$$+ \sum_{q=1}^{n_r} (-n_r)_q \frac{B_{q+1}}{(q+1)!} \zeta_{r-1}(-n_1, \dots, -n_{r-2}, -n_{r-1} - n_r + q).$$

Theorem (L.Jiu, V.H.Moll and C.Vignat, 2018)

$$\zeta_r(-n_1, \dots, -n_r) = \prod_{k=1}^r (-1)^{n_k} \mathcal{C}_{1, \dots, k}^{n_k+1}$$

Theorem (L.Jiu, V.H.Moll and C.Vignat, 2018)


$$\zeta_r(-n_1, \dots, -n_r) = \prod_{k=1}^r (-1)^{n_k} \mathcal{C}_{1, \dots, k}^{n_k+1},$$

$$\mathcal{C}_1^n = \frac{\mathcal{B}_1^n}{n}, \mathcal{C}_{1,2}^n = \frac{(\mathcal{C}_1 + \mathcal{B}_2)^n}{n}, \dots, \mathcal{C}_{1, \dots, k+1}^n = \frac{(\mathcal{C}_{1, \dots, k} + \mathcal{B}_{k+1})^n}{n}.$$

Theorem (L.Jiu, V.H.Moll and C.Vignat, 2018)

$$\zeta_r(-n_1, \dots, -n_r) = \prod_{k=1}^r (-1)^{n_k} \mathcal{C}_{1, \dots, k}^{n_k+1},$$

$$\mathcal{C}_1^n = \frac{\mathcal{B}_1^n}{n}, \mathcal{C}_{1,2}^n = \frac{(\mathcal{C}_1 + \mathcal{B}_2)^n}{n}, \dots, \mathcal{C}_{1, \dots, k+1}^n = \frac{(\mathcal{C}_{1, \dots, k} + \mathcal{B}_{k+1})^n}{n}.$$


$$\begin{aligned} \zeta_2(-n, 0) &= (-1)^n \mathcal{C}_1^{n+1} \cdot (-1)^0 \mathcal{C}_{1,2}^{0+1} \\ &= (-1)^n \frac{\mathcal{C}_1 + \mathcal{B}_2}{1} \cdot \mathcal{C}_1^{n+1} \\ &= (-1)^n (\mathcal{C}_1^{n+2} + \mathcal{B}_2 \mathcal{C}_1^{n+1}) \\ &= (-1)^n \left(\frac{\mathcal{B}_1^{n+2}}{n+2} + \mathcal{B}_2 \frac{\mathcal{B}_1^{n+1}}{n+1} \right) \\ &= (-1)^n \left[\frac{\mathcal{B}_{n+2}}{n+2} - \frac{1}{2} \frac{\mathcal{B}_{n+1}}{n+1} \right] \end{aligned}$$

Theorem (L.Jiu, V.H.Moll and C.Vignat, 2018)

$$\zeta_r(-n_1, \dots, -n_r) = \prod_{k=1}^r (-1)^{n_k} \mathcal{C}_{1, \dots, k}^{n_k+1},$$

$$\mathcal{C}_1^n = \frac{\mathcal{B}_1^n}{n}, \mathcal{C}_{1,2}^n = \frac{(\mathcal{C}_1 + \mathcal{B}_2)^n}{n}, \dots, \mathcal{C}_{1, \dots, k+1}^n = \frac{(\mathcal{C}_{1, \dots, k} + \mathcal{B}_{k+1})^n}{n}.$$

$$\begin{aligned} \zeta_2(-n, 0) &= (-1)^n \mathcal{C}_1^{n+1} \cdot (-1)^0 \mathcal{C}_{1,2}^{0+1} \\ &= (-1)^n \frac{\mathcal{C}_1 + \mathcal{B}_2}{1} \cdot \mathcal{C}_1^{n+1} \\ &= (-1)^n (\mathcal{C}_1^{n+2} + \mathcal{B}_2 \mathcal{C}_1^{n+1}) \\ &= (-1)^n \left(\frac{\mathcal{B}_1^{n+2}}{n+2} + \mathcal{B}_2 \frac{\mathcal{B}_1^{n+1}}{n+1} \right) \\ &= (-1)^n \left[\frac{\mathcal{B}_{n+2}}{n+2} - \frac{1}{2} \frac{\mathcal{B}_{n+1}}{n+1} \right]. \end{aligned}$$

$$\zeta(-n) = (-1)^n \mathcal{C}^{n+1} = (-1)^n \frac{\mathcal{B}_{n+1}}{n+1}.$$


Lattice/Motzkin Path

$$(m_n)_{n=0}^{\infty} \sim m_n = \int_{\mathbb{R}} x^n d\mu(x) \stackrel{?}{\Rightarrow} (P_n(x))_{n=1}^{\infty} \sim \int_{\mathbb{R}} P_n(x) P_m(x) d\mu(x) = C_n \delta_{m,n}$$

$$\Rightarrow P_{n+1}(x) = (x + s_n) P_n(x) + t_n P_{n-1}(x)$$

$$\Rightarrow \sum_{n=0}^{\infty} m_n x^n = \frac{m_0}{1 + s_0 x + \frac{t_1 x^2}{1 + s_1 x + \frac{t_2 x^2}{\ddots}}}$$

$$M_{n+1,k} = M_{n,k-1} + s_k M_{n,k} + t_{k+1} M_{n,k+1}$$



$$\sum_{n=0}^{\infty} M_{n,0} z^n = \frac{1}{1 - s_0 z - \frac{t_1 z^2}{1 - s_1 z - \frac{t_2 z^2}{\ddots}}}$$

Main Results: $H_n(a_k)$ for the following sequences

$$B_{2k+1} \left(\frac{x+1}{2} \right), E_{2k} \left(\frac{x+1}{2} \right), E_{2k+1} \left(\frac{x+1}{2} \right), E_{2k+2} \left(\frac{x+1}{2} \right),$$

$$B_k \left(\frac{x+r}{q} \right) - B_k \left(\frac{x+s}{q} \right), E_k \left(\frac{x+r}{q} \right) \pm E_k \left(\frac{x+s}{q} \right),$$


$$kE_{k-1}(x), B_{k+1, \times 8, 1}(x), B_{k+1, \times 8, 2}(x), B_{k+1, \times 12, 1}(x), B_{k+1, \times 12, 2}(x),$$

$$(2k+1)E_{2k}, (2^{2k+2} - 1)B_{2k+2}, (2k+1)B_{2k} \left(\frac{1}{2} \right), (2k+3)B_{2k+2},$$

| | | | | | | |
|-----------------|-------------------------------|---|----------------|---------------|----------------------|---------------------|
| $a_k, k \geq 1$ | B_{k-1} | B_{2k} | $(2k+1)B_{2k}$ | | $(2^{2k} - 1)B_{2k}$ | |
| a_0 | 0 | 1 | 1 | | 0 | |
| $a_k, k \geq 1$ | E_{2k-2} | $E_{k-1}(1)$ | $E_{k+3}(1)$ | $E_{2k-1}(1)$ | $E_{2k+5}(1)$ | $\frac{E_k(1)}{k!}$ |
| a_0 | 0 | 0 | $-\frac{1}{4}$ | 0 | $\frac{1}{2}$ | 1 |
| $a_k, k \geq 1$ | $\frac{E_{2k-1}(1)}{(2k-1)!}$ | $E_{2k-2} \left(\frac{x+1}{2} \right)$ | $(2k+1)E_{2k}$ | | | |
| a_0 | 0 | 0 | 0 | | | |

$$\frac{B_{2k+1} \left(\frac{x+1}{2} \right)}{2k+1}, \frac{B_{2k+3} \left(\frac{x+1}{2} \right)}{2k+3}, \frac{B_{2k+5} \left(\frac{x+1}{2} \right)}{2k+5}$$


$$B_{n+1} \left(\frac{x+2}{5} \right) - B_{n+1} \left(\frac{x}{5} \right) = \frac{n+1}{5^n} \sum_{k=0}^{\infty} \frac{3^k}{2^{2k+3}} \sum_{\ell=0}^k \binom{k}{\ell} (-1)^\ell \frac{1}{12^\ell} E_n^{(2k+2\ell+3)} (k+\ell+x).$$



Theorem (L. Jiu, I. Simonelli, and H. Yue, 21'+)

$$\phi_{0 \rightarrow a_n} = \phi_{0 \rightarrow a_1} \phi_{a_1 \rightarrow a_2 | \phi} \cdots \phi_{a_{n-1} \rightarrow a_n | a_{n-2}} \cdot \frac{1}{1 - P(L_1, \dots, L_n)},$$

$$B_{n+1} \left(\frac{x+2}{5} \right) - B_{n+1} \left(\frac{x}{5} \right) = \frac{n+1}{5^n} \sum_{k=0}^{\infty} \frac{3^k}{2^{2k+3}} \sum_{\ell=0}^k \binom{k}{\ell} (-1)^\ell \frac{1}{12^\ell} E_n^{(2k+2\ell+3)} (k+\ell+x).$$



Theorem (L. Jiu, I. Simonelli, and H. Yue, 21'+)


$$\phi_{0 \rightarrow a_n} = \phi_{0 \rightarrow a_1} \phi_{a_1 \rightarrow a_2} \cancel{\phi} \cdots \phi_{a_{n-1} \rightarrow a_n} \cancel{a_n \rightarrow 2} \cdot \frac{1}{1 - P(L_1, \dots, L_n)},$$

where

$$L_j = \phi_{a_{j-1} \rightarrow a_j} \cancel{a_j \rightarrow 2} \phi_{a_j \rightarrow a_{j-1}} \cancel{a_{j+1} \rightarrow 1}$$

$$P(L_1, \dots, L_n) = \sum_{*} (-1)^{\ell+1} L_{j_1} \cdots L_{j_\ell},$$

$$B_{n+1} \left(\frac{x+2}{5} \right) - B_{n+1} \left(\frac{x}{5} \right) = \frac{n+1}{5^n} \sum_{k=0}^{\infty} \frac{3^k}{2^{2k+3}} \sum_{\ell=0}^k \binom{k}{\ell} (-1)^\ell \frac{1}{12^\ell} E_n^{(2k+2\ell+3)} (k+\ell+x).$$



Theorem (L. Jiu, I. Simonelli, and H. Yue, 21'+)

$$\phi_{0 \rightarrow a_n} = \phi_{0 \rightarrow a_1} \phi_{a_1 \rightarrow a_2 | \phi} \cdots \phi_{a_{n-1} \rightarrow a_n | a_{n-2}} \cdot \frac{1}{1 - P(L_1, \dots, L_n)},$$

where

$$L_j = \phi_{a_{j-1} \rightarrow a_j | a_{j-2}} \phi_{a_j \rightarrow a_{j-1} | a_{j+1}}$$

$$P(L_1, \dots, L_n) = \sum_{*} (-1)^{\ell+1} L_{j_1} \cdots L_{j_\ell},$$

for the condition * given by (1) $\ell = 1, 2, \dots, n$; (2) and $j_1 < j_2 - 1, j_2 < j_3 - 1, \dots, j_{\ell-1} < j_\ell - 1$.

Future Work

- ▶ To apply this symbolic expression to the Tornheim zeta function, defined as

$$\mathcal{W}(r, s, t) := \sum_{m, n \geq 1} \frac{1}{m^r n^s (m+n)^t},$$

and its multi-dimensional extension.

- ▶ Another type of zeta function to which we may apply the Bernoulli symbol is the hypergeometric zeta function $\zeta_{a,b}^H(s)$. Let $(z_{k;a,b})_{k=1}^{\infty}$ be the sequence of complex zeros of ${}_1F_1\left(\begin{smallmatrix} a \\ a+b \end{smallmatrix} \middle| t\right)$, defined by

$$\zeta_{a,b}^H(s) := \sum_{k \geq 1} z_{k;a,b}^{-s} \quad \text{for } \text{Res} > 1.$$

- ▶ Orthogonal polynomials w. r. t. $B_n^{(p)}(x)$.
- ▶ Computer proofs for identities involving Bernoulli and Euler polynomials
- ▶ Hyperbolic secant (square) distribution and information geometry
- ▶

Future Work

- ▶ To apply this symbolic expression to the Tornheim zeta function, defined as

$$\mathcal{W}(r, s, t) := \sum_{m, n \geq 1} \frac{1}{m^r n^s (m+n)^t},$$

and its multi-dimensional extension.

- ▶ Another type of zeta function to which we may apply the Bernoulli symbol is the hypergeometric zeta function $\zeta_{a,b}^H(s)$. Let $(z_{k;a,b})_{k=1}^{\infty}$ be the sequence of complex zeros of ${}_1F_1\left(\begin{smallmatrix} a \\ a+b \end{smallmatrix} \middle| t\right)$, defined by

$$\zeta_{a,b}^H(s) := \sum_{k \geq 1} z_{k;a,b}^{-s} \quad \text{for } \text{Res} > 1.$$

- ▶ Orthogonal polynomials w. r. t. $B_n^{(p)}(x)$.
- ▶ Computer proofs for identities involving Bernoulli and Euler polynomials
- ▶ Hyperbolic secant (square) distribution and information geometry
- ▶

Thank you!!!